

NLO PARTONIC EVENT GENERATORS

S. WEINZIERL

*Dipartimento di Fisica, Università di Parma,
INFN Gruppo Collegato di Parma, 43100 Parma, Italy*



Merging the benefits of an NLO calculation with event generators is a topic of current interest. Among other things, it is advantageous to be able to generate unweighted events according to NLO matrix elements. I report on an algorithm which generates a sequence of unweighted momentum configurations, each configuration containing either n or $n+1$ four-vectors, such that for any infrared safe observable the average over these configurations coincides with the NLO calculation up to errors of a resolution variable.

1 Introduction

As high energy experiments move towards a new generation of colliders, accurate simulation of collision events becomes increasingly important. In particular this is the case for QCD, which often constitutes the bulk of the events. Theoretical predictions are in most cases either based on exact perturbative calculations, resummations or event generators. These methods have complementary regions of applicability. Exact perturbative calculations give a prediction for an observable at the parton level as a power series in the coupling constant. The state-of-the-art are numerical next-to-leading order programs, which allow the calculation of any infra-red safe observable for a given process. They give a reliable prediction, if all scales entering an observable are of the same order of magnitude, but are limited to a few partons only. If an observable depends on a large ratio of two scales, large logarithms appear in the perturbative expansion. In this situation one often reorganises the perturbative expansion and resums the large logarithms to all orders. Within resummation, the state of the art is next-to-leading logarithmic accuracy. Up to now, these resummations have to be carried out for each observable individually. Event generators follow a different philosophy and rely on a three-stage process: Starting from a hard-scattering process, parton showers are generated in the second stage. Finally, a hadronization model converts partons into hadrons. The first two stages, hard scattering and parton showering,

can be described in perturbation theory, while hadronization involves long-distance physics and is non-perturbative. Event generators usually only employ leading-order matrix elements and parton showers accurate to leading logarithmic accuracy. They describe reasonably well multiple collinear emission and they generate unweighted events, since they only use leading-order matrix elements. However, they do a rather poor job for multi-jet events. This is due to the fact that they only use the simplest $2 \rightarrow 2$ and $2 \rightarrow 3$ LO matrix elements, that the showering algorithm does not populate the complete phase space and that color interference is only approximated by angular ordering. Since multi-jet events become increasingly important for new physics searches, it is a topic of current interest to merge the benefits of exact perturbative calculations and event generators. Several issues have to be addressed for an event generator according to NLO matrix elements:

- Double counting: Gluon emission can originate from the hard scattering as well as from parton showering.
- Color flow: Matrix elements are usually color summed, where as the algorithms for showering and hadronization are based on open color lines.
- Negative weights: A standard NLO program generates n - and $(n+1)$ -parton phase space contributions, which are finite, but not necessarily positive definite.

Recent developments in this field considered combining parton showers with multi-parton tree-level matrix elements^{1–3}, with resummations of soft gluons⁴, and with NLO matrix elements^{5–9}. In this talk I focus on the issue of negative weights. Negative weights are not a problem per se, but can seriously waste computer resources, if events are passed through a detector simulation. It is generally the case that the CPU time needed for detector simulation exceeds the one needed for the generation of an event. It is therefore desirable to avoid them from the very beginning. In standard NLO calculations negative weights occur since individual contributions to an observable are not necessarily positive definite. It is therefore natural, to consider first a “partonic event generator”, leaving parton showers and hadronization for the moment aside. Here I report on an algorithm which generates a sequence of unweighted momentum configurations, each configuration containing either n or $n+1$ four-vectors, such that for any infrared safe observable the average over these configurations coincides with the NLO calculation up to errors of a resolution variable¹⁰. I focus on electron-positron annihilation with massless partons. The algorithm is independent of the specific hard process and can be implemented on top of existing NLO programs. After an outline of the basic idea, I focus on a few technical challenges which have to be addressed in order for the algorithm to run efficiently.

2 The algorithm

Before stating the algorithm I briefly recall the main features of standard NLO calculations. The calculation of an observable at NLO receives contributions from the Born matrix element, the virtual corrections and the real emissions. Taken separately, the last two contributions are divergent, only their sum is finite. The subtraction- or phase space slicing method can be used to render them finite individually. Within the subtraction method one adds and subtracts a suitable piece $d\sigma^A$ ¹¹:

$$\sigma^{NLO} = \int_n d\sigma^B + \int_n \left(d\sigma^V + \int_1 d\sigma^A \right) + \int_{n+1} (d\sigma^R - d\sigma^A) \quad (1)$$

The modified virtual corrections and the modified real emissions are now separately finite, but not necessarily positive definite. The basic idea to generate unweighted events according to NLO

matrix elements is relatively simple and inspired by a slicing approach (although it will entirely be based on the subtraction method): Consider a $(n + 1)$ -parton configuration. If all partons are separated by at least y_{res} according to a resolution criteria, one returns this $(n + 1)$ -parton configuration. If on the other hand a parton is not resolved at a scale y_{res} , one averages over the real emissions up to the scale y_{res} , combines the result with the virtual corrections and returns a n -parton configuration. By choosing y_{res} large enough the sum will have a positive weight. For an efficient implementation I would like to discuss three technical points:

- A modified version of the subtraction method, such that the subtraction terms are non-vanishing only for $y < y_{res}$.
- A parameterization of the phase space, which starts from a n parton configuration and inserts an additional soft or collinear particle, such that the integration over the unresolved part involves only a three-dimensional integration and, in addition, such that the dominant $1/s_{ij}$ -behaviour of the integrand is absorbed into the measure. This ensures that the unresolved region can be integrated efficiently.
- The Metropolis algorithm to generate events with constant weight according to a (complicated) probability distribution.

I start with a modified version of the subtraction method: The subtraction method has in general the advantage that it does not introduce additional approximations to the matrix elements or to the phase space. The only approximation made for the algorithm presented here, consists in replacing a $(n + 1)$ -parton configuration by a n -parton configuration, whenever two partons are not resolved within y_{res} . In order for the basic idea outlined above to work, nothing should be subtracted if all partons are separated by y_{res} . In the original version of the dipole formalism¹¹ the approximation term is given by a sum over dipoles $\sum \mathcal{D}_{ij,k}$ and subtracted over the complete phase space. However, it turns out that the approximation term can still be integrated analytically if the region where it is subtracted, is restricted¹²:

$$d\sigma^A = \sum \mathcal{D}_{ij,k} \Theta(y_{ij,k} < y_{res}) \quad (2)$$

Here, $y_{ij,k} = 2p_i p_j / (2p_i p_j + 2p_i p_k + 2p_j p_k)$ is the appropriate resolution variable if partons i and j become close to each other and k is the spectator.

The second point is a suitable parametrization of the phase space. For any given “hard” n -parton configuration one needs a parametrization for the insertion of an additional parton, which can become soft or collinear to one of the n hard partons. This is done as follows¹³: The phase space measure of the $(n + 1)$ -parton configuration can be related to a n -parton configuration by

$$d\phi_{n+1} = d\phi_n \frac{ds_{as} ds_{sb} d\phi_s}{4(2\pi)^3 s_{ab}}. \quad (3)$$

The invariants s_{as} , s_{sb} and the angle ϕ_s are parametrized in terms of three random variables v_1 , v_2 and v_3 , uniformly distributed in $[0, 1]$:

$$s_{as} = s_{ab} \left(\frac{s_{min}}{s_{ab}} \right)^{v_1}, s_{sb} = s_{ab} \left(\frac{s_{min}}{s_{ab}} \right)^{v_2}, \phi_s = 2\pi v_3. \quad (4)$$

s_{min} is an internal technical parameter, which is chosen to be small. It is possible to reconstruct the four-vectors of the $n + 1$ parton configuration from these invariants. The phase space measure becomes

$$d\phi_{n+1} = d\phi_n \frac{1}{4(2\pi)^2} \frac{s_{as} s_{sb}}{s_{ab}} \ln^2 \left(\frac{s_{min}}{s_{ab}} \right) dv_1 dv_2 dv_3 \quad (5)$$

and the $1/s_{ij}$ -behaviour of the integrand has been absorbed into the measure.

With an appropriate choice of y_{res} one obtains now events with positive weights. In general, the probability distribution will be a rather complicated function and unweighting according to the standard acceptance-rejection method might not be feasible. The method of choice is therefore the Metropolis algorithm, which allows to generate unweighted events according to a multi-dimensional probability density $P(x_1, \dots, x_d)$, which not necessarily factorizes. One starts from a state $\vec{x} = (x_1, \dots, x_d)$ and randomly generates a new candidate $\vec{x}'$. One then calculates the quantity $\Delta S = -\ln(P(\vec{x}')/P(\vec{x}))$ and accepts the new candidate with probability $\min(1, e^{-\Delta S})$. The way new candidates are suggested is arbitrary, restricted only by ergodicity and detailed balance: Each state must be reachable within a finite number of steps and the probability of suggesting the state x_2 given we are in state x_1 has to be equal to the probability of suggesting x_1 given we are in x_2 .

To summarize, the complete algorithm to generate unweighted events according to NLO matrix elements is therefore as follows: One first picks a set of random numbers $u_1, \dots, u_m, x, v_1, v_2, v_3$, where $u_1, \dots, u_m$ define the “hard” n -parton configuration, x decides where the “soft” particle gets inserted and v_1, v_2, v_3 (together with the “hard” configuration and the information where the “soft” particle gets inserted) yield the “soft” $(n+1)$ event. One then checks for the $(n+1)$ -parton event if all $y_{ij,k}$ are greater than y_{res} , e.g. if all partons are well separated. If this is the case the probability P' is proportional to the real emission matrix element and the $(n+1)$ parton configuration will be returned, if the event is accepted by the Metropolis algorithm. If not all partons are well separated, the “hard” n -parton configuration has the probability

$$P' \sim |\mathcal{M}_{Born}^n|^2 + 2 \operatorname{Re} \mathcal{M}_{Born}^n * \mathcal{M}_{1-loop}^n + \langle \mathcal{M}_{Born}^n | \mathbf{I} | \mathcal{M}_{Born}^n \rangle + \int dv_1 dv_2 dv_3 w_{soft} \left(|\mathcal{M}_{Born}^{n+1}|^2 - \sum \mathcal{D}_{ij,k} \Theta(y_{ij,k} < y_{res}) \right) \quad (6)$$

The last terms in the first and second line are the added and subtracted dipole terms, which ensure that the expressions over the n - and $(n+1)$ -parton phase space are finite separately. The second line involves a three-dimensional integration over the unresolved phase space. In both cases, e.g. if the event is classified as a n - or $(n+1)$ -parton event, the new candidate gets accepted with Metropolis probability $\min(1, P'/P)$, where P' is the probability of the new candidate and P is the probability of the previous event.

3 Outlook

I presented an algorithm to generate unweighted partonic events according to NLO matrix elements in electron-positron annihilation. An important ingredient is an efficient integration over the unresolved region on a point-by-point basis in hard phase space. The extension to initial-state partons or massive partons should be feasible.

References

1. M. H. Seymour. *Comp. Phys. Commun.*, 90:95–101, 1995.
G. Corcella and M. H. Seymour. *Phys. Lett.*, B442:417–426, 1998.
2. M. L. Mangano, M. Moretti, and R. Pittau. hep-ph/0108069.
3. S. Catani, F. Krauss, R. Kuhn, and B. R. Webber. *JHEP*, 11:063, 2001.
4. S. Mrenna. hep-ph/9902471.
5. C. Friberg and T. Sjostrand. hep-ph/9906316.
6. B. Potter. *Phys. Rev.*, D63:114017, 2001.
B. Potter and T. Schorner. *Phys. Lett.*, B517:86–92, 2001.

7. M. Dobbs and M. Lefebvre. *Phys. Rev.*, D63:053011, 2001.
M. Dobbs. *Phys. Rev.*, D64:034016, 2001.
M. Dobbs. *Phys. Rev.*, D65:094011, 2002.
8. S. Frixione and B. R. Webber. hep-ph/0204244.
9. J. C. Collins. *JHEP*, 05:004, 2000.
J. C. Collins and F. Hautmann. *JHEP*, 03:016, 2001.
Y. Chen, J. C. Collins, and N. Tkachuk. *JHEP*, 06:015, 2001.
J. C. Collins. hep-ph/0110113.
J. C. Collins and X. -M. Zu. hep-ph/0204127.
10. S. Weinzierl. *JHEP*, 08:028, 2001.
11. S. Catani and M. H. Seymour. *Nucl. Phys.*, B485:291–419, 1997.
12. Z. Nagy and Z. Trocsanyi. *Phys. Rev.*, D59:014020, 1999.
13. S. Weinzierl and D. A. Kosower. *Phys. Rev.*, D60:054028, 1999.